CONTENTS

The inversion of correlation between γ-effect values and dihedral angle: nontraditional chair–boat conformational equilibrium in seven- and eight-membered dithioacetals
E.N. Klimovitskii,I I.A. Litvinov, O.N. Kataeva, D.Yu. Strel'nik and G.N. Sergeeva (Kazan, U.S.S.R.) 1

The synthesis and crystal structures of three Mo–Cu–S cubane-like clusters: $[Mo_3CuS_4][S_2P(OCH_2CH_3)_2]_3 \cdot I \cdot (\mu_2\text{-}OOCCH3) \cdot H_2O$, $[Mo_3CuS_4][S_2P(OCH_2CH_3)_2]_3 \cdot I \cdot (\mu_2\text{-}OOCCH_3) \cdot (CH_3)_2SO$ and $[Mo_3CuS_4][S_2P(OCH_2CH_3)_2]_3 \cdot I \cdot (\mu_2\text{-}OOCC_6H_5 \cdot C_5H_5N$
S.-F. Lu, N.-Y. Zhu, X.-T. Wu, Q.-J. Wu and J.-X. Lu (Fujian, People's Republic of China) ... 15

The synthesis and crystal structure of a novel cubane-like tungsten copper sulfur cluster $[W_3CuS_4] \cdot [S_2p(OC_2H_5)_2]_3 \cdot I \cdot \mu_2\text{-}CH_3COO \cdot CH_3CN$
X.-T. Wu, H.-Q. Zhan, Y.-F. Zheng and J.-X. Lu (Fujian, People's Republic of China) 33

Crystal and molecular structure of the MEGA-SPIRO cyclophosphazenic species from diethylene glycol bis-(3-aminopropyl)-ether, $N_3P_3Cl_4\{(HN–(CH_2)_3–O–(CH_2)_2–O–(CH_2)_2–O–(CH_2)_3–NH)\}$
T.S. Cameron and A. Linden (Halifax, N.S., Canada), F. Sournies, A. El Bakili and J.-F. Labarre (Toulouse, France) 41

NQR study of complexes between *p*-dichlorobenzene and some aromatic π acceptors
N. Subramani and K.V. Raman (Tiruchirapalli, India) and J. Ramakrishna (Bangalore, India) 53

Synthesis and structural study of tropane benzamines. Part III. *N*-(8-Isopropyl-nortropan-3-β-yl)-2-methoxy-4-amino-5-chlorobenzamide
N. Cabezas, M. Martinez, E. Galvez, M.S. Arias, F. Florencio and J. Sanz-Aparicio (Madrid, Spain) 59

Some effects of intramolecular hydrogen bonding on vibrational spectra
V.M. Schreiber (Leningrad, U.S.S.R.) 73

Vibrational spectrum and internal rotation in 2,6-dimethylpyrazine
J.F. Arenas, J.T. Lopez-Navarrete, J.I. Marcos and J.C. Otero (Malaga, Spain) 87

NH stretching vibrations and conformation of bis[2-(3-substituted ureido)phenyl] disulphides
A.Ts. Antonova (Sofia, Bulgaria) 97

Infrared, Raman and pre-resonance Raman spectroscopy of $Rh_2(O_2CCH_3)_4(S(CH_2Ph)_2)_2$
R.J.H. Clark and A.J. Hempleman (London, Gt. Britain) 105

Incoherent inelastic neutron scattering studies of protonconducting materials: $Sn(HPO_4)_2 \cdot H_2O$ and $HM(SO_4)_2 \cdot H_2O$, M=Fe, In. Part II. The vibrational spectrum of H_3O^+
D.J. Jones (Montpellier, France), J. Penfold and J. Tomkinson (Didcot, Gt. Britain) and J. Rozière (Montpellier, France) 113

Additional evidence concerning the conformational equilibrium of 2-aminoethanethiol from the microwave spectrum of the ND_2–SD trideuterated species and ab initio calculations
W. Caminati and B. Belino (Bologna, Italy), L. Schäfer and J.D. Ewbank (Fayetteville, AR, U.S.A.) and K. Siam (Pittsburg, KS, U.S.A.) 123

Conformational analysis of 1,2-dihalotetramethylethanes and 1,2-dihalotetramethyldisilanes by molecular mechanics calculations
R. Stølevik and P. Bakken (Dragvoll, Norway) 131

Conformational analysis of *cis, cis*-2,2′- and 1,1′-halogensubstituted bicyclopropyl derivatives by molecular mechanics calculations
R. Stølevik and P. Bakken (Dragvoll, Norway) 137

Far-infrared spectra and conformational stability of the epihalohydrins
J.R. Durig, S.E. Godbey and R.A. Larsen (Columbia, SC, U.S.A.) 143

Thiocarbonyl spectroscopy: the infrared spectrum and ab initio vibrational frequencies of *cis*- and *trans*-dithioformic acid in the X^1A' ground state
F. Ioannoni and D.C. Moule (St. Catharines, Ont., Canada), J.D. Goddard (Guelph, Ont., Canada) and D.J. Clouthier (Lexington, KY, U.S.A.) 159

Raman spectroscopic study of 1,2-ethanedithiol adsorbed on silver surface
C.K. Kwon, K. Kim, M.S. Kim and Y.S. Lee (Seoul, Korea) 171

Ground-state rotational spectrum of t-butyl isocyanide and an r_s geometry of its heavy-atom skeleton
N.W. Howard, A.C. Legon, C.A. Rego and A.L. Wallwork (Exeter, Gt. Britain) 181
Spectroscopic study of the conformational isomerism of 2-formylfuran derivatives
A.I. Kiss and D. Machytka (Budapest, Hungary), J. Bánki (Bratislava, Czechoslovakia) and M. Gál (Budapest, Hungary) 193
The vibrational spectrum of trimethylphosphine–silver iodide, $[AgI{\cdot}P(CH_3)_3]_4$
H.G.M. Edwards and D.W. Farwell (Bradford, Gt. Britain) 203
A transferable general valence force field for oxalyl halides: Fourier transform infrared spectral analysis of oxalyl chloride isolated in cryogenic rare gas matrices
G. Davidovics, M. Monnier, W. Schroeder, P. Verlaque, J. Pourcin and H. Bodot (Marseille, France) 213
Ultraviolet photoisomerizations and FT-IR investigations of matrix isolated oxalyl halide conformers
W. Schroeder, M. Monnier, G. Davidovics, A. Allouche, P. Verlaque, J. Pourcin and H. Bodot (Marseille, France) 227
Dielectric relaxation mechanism and internal rotation in phenylpyridines
H.A. Kołodziej and E. Harewski (Wrocław, Poland) 243
The relation of methoxy group CH bond lengths and stretching frequencies to the oxygen lone pair orientation
I.S. Ignatyev (Leningrad, U.S.S.R.) 251
Contributions to the chemistry of silicon–sulphur compounds. Part 54. Extension of the molecular mechanics parameterization to polyheteroatomic compounds
Z. Pawelec, A. Herman and W. Wojnowski (Gdansk, Poland) 259
Electronic and geometric structures of oligothiophenes
U. Nagashima, H. Fujimoto and H. Inokuchi (Okazaki, Japan) and K. Seki (Hiroshima, Japan) 265
Photoelectron spectra and electronic structure of some 1-substituted aziridines and of 1,1′-biaziridinyl
P. Rademacher, G. Irsch and W. Sicking (Essen, F.R.G.) and E.-U. Würthwein (Münster, F.R.G.) 291
Application of Schwinger perturbation theory in electron diffraction analysis. Part I. Linear XY_2 type molecules
K.V. Ermakov, B.S. Butayev and V.P. Spiridonov (Moscow, U.S.S.R.) 307
The gas-phases structures of *trans*- and *cis*-perfluorodecalin
H.-G. Mack and H. Oberhammer (Tübingen, F.R.G.) 321
The gas-phase structures of perfluoroquinolizidine and perfluorotripropylamine
H.-G. Mack and H. Oberhammer (Tübingen, F.R.G.) 329
The molecular structure and conformational equilibrium of 1,2-diisocyano-ethane studied by gas-phase electron diffraction
G. Schrumpf (Göttingen, F.R.G.), M. Traetteberg and P. Bakken (Dragvoll, Norway) and R. Seip (Oslo, Norway) 339
The molecular structure of trimethylantimony(V)dichloride as determined by gas-phase electron diffraction
Q. Shen (Hamilton, NY, U.S.A.) and R.T. Hemmings (Mississauga, Ont., Canada) 349

Short communication
An infrared and Raman spectroscopic study of transition metal(II)chloride 2,6-lutidine complexes
S. Yurdakul (Ankara, Turkey), S. Akyuz (Samsun, Turkey) and J.E.D. Davies (Lancaster, Gt. Britain) 355
Calculations of induced dipole moments: adamantane derivatives
L. Došen-Mićović (Belgrade, Yugoslavia) and O. Exner (Prague, Czechoslovakia) 361
Conformational analysis of 2-heptyne
G.A. Crowder (Canyon, TX, U.S.A.) 367

Book review 373
Author index 375
Subject index 376